

Cabenuva

(cabotegravir; rilpivirine)

Provider Order Form rev. 2/9/2025



PROVIDERS: Please include the following information to expedite the order:
Patient Demographic, most recent Office Visit Note, Insurance Information, TB Results, Recent Labs

PATIENT INFORMATION

Referral Status: New Referral Updated Order Order Renewal

Date: _____ Patient Name: _____ DOB: _____
ICD-10 code: B20 Other: _____
Description: _____
 NKDA Allergies: _____ Weight (lb/kg) _____
Patient Status: New to Therapy New to Therapy Last Treatment Date: _____ Next Due Date: _____

PROVIDER INFORMATION

Referral Coordinator Name: _____ Referral Coordinator Email: _____
Ordering Provider: _____ Provider NPI: _____
Referring Practice Name: _____ Phone: _____ Fax: _____
Practice Address: _____ City: _____ State: _____ Zip: _____

NURSING

- TB status and date (list results and attach clinicals)

- Utilize Hypersensitivity protocol established by Redeemer Health Infusion and PI
- Provide nursing care and observation per Redeemer Health Infusion Policies and Procedures

LABORATORY ORDERS

- CBC at each dose every _____
- CMP at each dose every _____
- CRP at each dose every _____
- Other: _____

Please check this box if you do NOT authorize Redeemer Health Infusion to order and draw labs indicated for clinical clearance and/or insurance authorization prior to treatment.

PRE-MEDICATION ORDERS

Not usually indicated for this medication.

- acetaminophen (Tylenol) 500 mg/ 650 mg/ 1000 mg PO
- cetirizine (Zyrtec) 10 mg PO
- loratadine (Claritin) 10 mg PO
- diphenhydramine (Benadryl) 25 mg/ 50 mg PO/ IV
- methylprednisolone (Solu-Medrol) 40 mg/ 125 mg IV
- hydrocortisone (Solu-Cortef) 100 mg IV
- Other: _____
Dose: _____ Route: _____
Frequency: _____

THERAPY ADMINISTRATION

- Cabenuva Intramuscular Injection
 - **Monthly Dosing:**
 - Initiation: 600/900 mg on last day of current antiretroviral therapy or oral lead-in
 - Maintenance 400mg/600 mg every month
 - **Every 2-Month Dosing:**
 - Initiation: 600 mg/900 mg on last day of current antiretroviral therapy or oral lead-in and 600 mg/900 mg again one month later.
 - Maintenance: 600 mg/900 mg two months after second initiation dose and every two months thereafter
 - Other: _____
- Doses may be given +/- 7 days from date of scheduled regimen.
- Route:** Intramuscular
- Refills: Zero/ for 12 months/ Other: _____
(If not indicated, order will expire one year from date signed)

To ensure a brand name product is dispensed, the prescriber must *handwrite "Brand Medically Necessary" on the prescription form. If not indicated, Redeemer Infusion is authorized to administer a generic or biosimilar.*

SPECIAL INSTRUCTIONS

Please check this box if you DO NOT authorize Redeemer Health Infusion to complete a Peer to Peer on behalf of the prescribing provider for an insurance company that denies authorization for treatment.

Provider Name (Print)

Provider Signature

Date