

Cabotegravir/Rilpivirine (CABENUVA)

PROVIDERS: Please include the following information to expedite the order:
Patient demographics, insurance information, most recent visit notes, and HIV-1 lab results

PATIENT INFORMATION

Patient: _____ DOB: _____
ICD-10: B20.____ Z21.____ Other: _____ Description: _____
Allergies: _____ NKDA Weight: _____ lb kg
Patient Status: New to Therapy Continuation of Therapy Last Treatment Date: _____ Next Due Date: _____

REFERRAL STATUS New Referral Updated Order Order Renewal

PROVIDER INFORMATION

Referral Coordinator Name: _____ Referral Coordinator Email: _____
Ordering Provider: _____ Provider NPI: _____
Referring Practice Name: _____ Phone: _____ Fax: _____
Practice Address: _____ City: _____ State: _____ Zip: _____

NURSING

- HIV-1 Lab Results (list result and attach documentation)

- Utilize hypersensitivity protocol established by Redeemer Health Infusion Therapy Center
- Provide nursing care and observation according to Redeemer Health Infusion Therapy Center policies and procedures

LABORATORY ORDERS

PRE-MEDICATION ORDERS

SPECIAL INSTRUCTIONS

THERAPY ADMINISTRATION

- Cabotegravir/Rilpivirine (CABENUVA) intramuscular injection

Monthly Dosing:

- Initiation: 600 mg/900 mg on the last day of current antiretroviral therapy or oral lead-in
- Continuation: 400 mg/600 mg every month (beginning one month after initiation dose)

Bimonthly Dosing:

- Initiation: 600 mg/900 mg on the last day of current antiretroviral therapy or oral lead-in for 2 consecutive months
- Continuation: 600 mg/900 mg every two months (beginning two months after second initiation dose)

Refills: No Refills 12 months _____

Redeemer Health Infusion Therapy Center will conduct peer-to-peer review(s) on behalf of the prescribing provider for any insurance denials. If you DO NOT AUTHORIZE Redeemer Health Infusion Therapy Center to do this on your behalf, check this box.

Provider Name (Print): _____ Provider Signature (No Stamps): _____ Date: _____