

Ocrelizumab (OCREVUS)

PROVIDERS: Please include the following information to expedite the order:
Patient demographics, insurance information, most recent visit notes, Hep B status, and applicable labs

PATIENT INFORMATION

Patient: _____ DOB: _____
ICD-10 (Required): _____ Description: _____
Allergies: _____ NKDA Weight: _____ lb kg
Patient Status: New to Therapy Continuation of Therapy Last Treatment Date: _____ Next Due Date: _____

REFERRAL STATUS New Referral Updated Order Order Renewal

PROVIDER INFORMATION

Referral Coordinator Name: _____ Referral Coordinator Email: _____
Ordering Provider: _____ Provider NPI: _____
Referring Practice Name: _____ Phone: _____ Fax: _____
Practice Address: _____ City: _____ State: _____ Zip: _____

NURSING

- Utilize hypersensitivity protocol established by Redeemer Health Infusion Therapy Center
- Provide nursing care and observation according to Redeemer Health Infusion Therapy Center policies and procedures
- Hepatitis B status and date (list results here and attach clinicals): _____
- Recent quantitative serum immunoglobulin test (list results here and attach clinicals): _____

LABORATORY ORDERS

- CBC at each dose every _____
- CMP at each dose every _____
- CRP at each dose every _____
- Treatment naïve patient required labs:** ALT, AST, Alkaline Phosphate, Bilirubin (list results here and attach clinicals): _____
- Please check this box if you do NOT authorize Redeemer Health Infusion Therapy Center to order and draw labs indicated for clinical clearance and/or insurance authorization prior to treatment.

PRE-MEDICATION ORDERS

- acetaminophen (TYLENOL) 500 mg 650 mg 1000 mg PO
- cetirizine (ZYRTEC) 10 mg PO
- loratadine (CLARITIN) 10 mg PO
- diphenhydramine (BENADRYL) 25 mg 50 mg | PO IV
- methylprednisolone (SOLU-MEDROL) 40 mg IV 125 mg IV
- hydrocortisone (SOLU-CORTEF) 100 mg IV

THERAPY ADMINISTRATION

- Ocrelizumab (OCREVUS)

Induction:

Dose: 300 mg in 250 mL 0.9% sodium chloride

Frequency: on day 1 and day 15

Rate: minimum of 2.5 hours according to package insert

Maintenance:

Dose: 600 mg in 500 mL 0.9% sodium chloride

Frequency: every 6 months from infusion 1 of initial dose

Rate: minimum of 2 hours according to package insert

Route: intravenous infusion

Flush: with 0.9% sodium chloride after infusion complete

Monitoring: Patient required to stay for 60-minute observation

Refills: No Refills 12 months _____

SPECIAL INSTRUCTIONS

Redeemer Health Infusion Therapy Center will conduct peer-to-peer review(s) on behalf of the prescribing provider for any insurance denials. If you DO NOT AUTHORIZE Redeemer Health Infusion Therapy Center to do this on your behalf, check this box.

Note to prescriber: Hepatitis B virus and quantitative serum immunoglobulin screening are required before the first dose. Pre-medicate with methylprednisolone (or an equivalent corticosteroid) and an antihistamine (e.g., diphenhydramine) prior to each infusion.

Provider Name (Print): _____ Provider Signature (No Stamps): _____ Date: _____